

Restoration of chiral symmetry in light-front finite temperature field theory*

S. Strauß**, M. Beyer, S. Mattiello

Fachbereich Physik, Universität Rostock, D-18051 Rostock, Germany

Abstract. We investigate the properties of qq and $q\bar{q}$ states in hot and dense quark matter in the framework of light-front finite temperature field theory. Presently we use the Nambu Jona-Lasinio model of QCD and derive the gap equation at finite temperature and density. We study pionic and scalar diquark dynamics in quark matter and calculate the masses and the Mott dissociation as a function of the temperature T and the chemical potential μ . For the scalar diquark we determine the critical temperature of color superconductivity.

1 In-medium gap equation in NJL

Relativistic heavy ion collisions open the possibility to study quantum chromodynamics (QCD) at finite temperature T and density or chemical potential μ . This is necessary to understand the evolution of the early universe and the interior of neutron stars. A way suited to describe the QCD phases is using light-front quantization. Previously we have used a spinless zero-range interaction. Here we use the light-front form [1] of the Nambu Jona-Lasinio (NJL) model that shows chiral symmetry breaking and calculate several phase boundaries including the chiral symmetry restoration. The NJL Lagrangian for the two flavor case reads

$$\mathcal{L}_{\text{NJL}} = \bar{\psi}(i\gamma_\mu\partial^\mu - m_0)\psi + G((\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma_5\boldsymbol{\tau}\psi)^2). \quad (1)$$

An extensive discussion on the NJL model, the chiral symmetry breaking, and the properties of light mesons at finite temperatures and chemical potential in the instant form can be found in Ref. [2].

On the light-front the in-medium quark propagator [3] is given as

$$\mathcal{G}(k) = (\gamma_\mu k_\mu^\mu + m) \left\{ \frac{1 - f^+(k^+, \mathbf{k}_\perp)}{k^2 - m^2 + i\varepsilon} + \frac{f^+(k^+, \mathbf{k}_\perp)}{k^2 - m^2 - i\varepsilon} \right\} \quad (2)$$

*Presented by S. Strauß at Light-Cone 2004, Amsterdam, 16 - 20 August

**email: strauss@rubin.physik2.uni-rostock.de

where $k_{\text{on}}^2 = m^2$ denotes the on-shell 4-momentum. The Fermi functions for a quark (f^+) or antiquark (f^-) are

$$f^\pm(k^+, \mathbf{k}_\perp) = \left[\exp \left\{ \frac{1}{T} \left(\frac{1}{2} k_{\text{on}}^- + \frac{1}{2} k^+ \mp \mu \right) \right\} + 1 \right]^{-1}. \quad (3)$$

For the generalization of the gap equation to finite T and μ it is most convenient to use the Dyson equation for the quark propagator and compute the self energy in Hartree approximation using (2). This leads to the in-medium gap equation

$$m(T, \mu) = m_0 + 24G \int_0^\infty \frac{dk^+}{k^+ (2\pi)^3} \int_{\mathbb{R}^2} d^2 \mathbf{k}_\perp m (1 - f^+(k^+, \mathbf{k}_\perp) - f^-(k^+, \mathbf{k}_\perp)). \quad (4)$$

We introduce the Lepage-Brodsky (LB) cut-off scheme to regulate the divergences in (4) and in the following.

2 Two-body states

The two-body bound states are investigated within the Bethe-Salpeter approach. We are using the following T -matrix equation [4]

$$T(k) = K + \int \frac{d^4 q}{(2\pi)^4} K S_F(q + k/2) S_F(q - k/2) T(k), \quad (5)$$

where K is an appropriate interaction kernel and $S_F(q)$ the dressed quark propagator. Bound states of mass M are identified with poles of the T -matrix $T(k)$ on the mass-shell $k^2 = M^2$.

2.1 Pion

The interaction in the pseudoscalar channel is given by

$$K_{\alpha\beta, \gamma\delta} = -2iG(\gamma_5 \tau_i)_{\alpha\beta} (\gamma_5 \tau_i)_{\gamma\delta}. \quad (6)$$

One introduces a reduced T -Matrix $t_\pi(k)$

$$T(k)_{\alpha\beta\gamma\delta} = (\gamma_5 \tau_i)_{\alpha\beta} t_\pi(k) (\gamma_5 \tau_i)_{\gamma\delta}. \quad (7)$$

Inserting the definition (7) into (5) the solution for $t_\pi(k)$ is obtained

$$t_\pi(k) = \frac{-2iG}{1 + 2G\Pi_\pi(k^2)} \quad (8)$$

with $\Pi_\pi(k^2)$ evaluated using (2) for $S_F(k)$

$$\Pi_\pi(k^2) = -6 \int_{LB} \frac{dx d^2 \mathbf{q}_\perp}{x(1-x)(2\pi)^3} \frac{M_{20}^2(x, \mathbf{q}_\perp) (1 - f^+(M_{20}) - f^-(M_{20}))}{M_{20}^2(x, \mathbf{q}_\perp) - k^2}. \quad (9)$$

where $M_{20}^2(x, \mathbf{q}_\perp) = (\mathbf{q}_\perp^2 + m^2)/x(1-x)$. For given T and μ one determines the value of k^2 to fulfill the pole condition $1 + 2G\Pi_\pi(m_\pi^2) = 0$. The results for the pion mass m_π are presented in Fig. 1.A.




Figure 1. (A) The pion mass as a function of T for different μ . (B) The diquark mass as a function of T for different μ . The continuum is given by $2m$. The lines of the two-body masses end at the Mott dissociation points.

2.2 Diquark

For the scalar, isospin singulet, color-antitriplet diquark the interaction kernel is

$$K_{\alpha\beta,\gamma\delta} = 2iG_s(\gamma_5 C \tau_2 \lambda^a)_{\alpha\beta} (C^{-1} \gamma_5 \tau_2 \lambda^a)_{\gamma\delta} \quad (10)$$

with color index $a = 2, 5, 7$ and $C = i\gamma^2\gamma^0$.

Following Ref. [4] one can construct the interaction in the diquark channel by a Fierz transformation of the NJL Lagrangian (1). We treat the coupling constant G_s as a free parameter to get a reasonably high diquark mass for $T = \mu = 0$. The solution of the T -matrix equation (5) with the kernel (10) is

$$T(k) = (\gamma_5 C \tau_2 \lambda^a)_{\alpha\beta} \tau_s(k) (C^{-1} \gamma_5 \tau_2 \lambda^a)_{\gamma\delta} \quad (11)$$

with

$$\tau_s(k) = \frac{2iG_s}{1 + 2G_s\Pi_s(k^2)} \quad (12)$$

and

$$\Pi_s(k^2) = -6 \int_{LB} \frac{dx d^2\mathbf{q}_\perp}{x(1-x)(2\pi)^3} \frac{M_{20}^2(x, \mathbf{q}_\perp) (1 - 2f^+(M_{20}))}{M_{20}^2(x, \mathbf{q}_\perp) - k^2}. \quad (13)$$

Results for the medium dependence of the diquark mass m_d are shown in Fig. 1.B.

3 Color superconductivity

Because of the attractive interaction between quarks one expects, by Cooper's theorem, a color superconducting phase (CSC) to be present for cold, dense quark matter. The critical temperature of CSC can be determined via the Thouless criterion that describes the condensation of the bosonic diquarks, viz.

$$m_d(T, \mu) = 2\mu. \quad (14)$$

Inserting relation (14) the pole condition $1 + 2G_s\Pi_s(m_d^2) = 0$ for the diquarks becomes

$$\frac{1}{2G_s} = 6 \int_{LB} \frac{dx}{x(1-x)} \int_{LB} \frac{d^2\mathbf{q}_\perp}{(2\pi)^3} \frac{M_{20}^2(x, \mathbf{q}_\perp) (1 - 2f^+(M_{20}))}{M_{20}^2(x, \mathbf{q}_\perp) - 4\mu^2}. \quad (15)$$




Figure 2. The phase diagram in the NJL model. The solid line shows the chiral phase transition. The transition between the deconfined and the CSC phase is given by the dotted line. And the dashed line is the Mott dissociation line of the pion.

Since (15) relates the critical temperature and chemical potential for CSC one can compute the phase transition from deconfined to color superconducting quark matter. The result is given in Fig. 2.

4 Conclusion

We have shown that the well known features of the NJL model (chiral phase transition, diquarks, pion dissociation, superconductivity) are recovered in a light-front quantization of statistical physics. This new approach to finite temperature field theory seems to work quite well. Despite all the challenges that might occur, when applied to real QCD, this opens a new perspective to explore QCD at finite temperatures and in particular at larger densities ($\mu > T$), where lattice QCD is facing severe technical problems.

Acknowledgment

We thank the organizers of the Light-Cone 2004 workshop in Amsterdam for this inspiring meeting. This work is supported by the Deutsche Forschungsgemeinschaft.

References

1. P.A.M. Dirac, Rev. Mod. Phys. **21**, 392 (1949)
2. S. P. Klevansky, Rev. Mod. Phys. **64** (1992) 649.
3. M. Beyer, S. Mattiello, T. Frederico and H. J. Weber, Phys. Lett. B **521** (2001) 33; M. Beyer, S. Mattiello, T. Frederico and H. J. Weber, arXiv:hep-ph/0310222.
4. N. Ishii, W. Bentz and K. Yazaki, Nucl. Phys. A **587** (1995) 617; W. Bentz, T. Hama, T. Matsuki and K. Yazaki, Nucl. Phys. A **651** (1999) 143.